

# IYGB- FP2 PAPER Q - QUESTION 1

START BY FINDING AN INTEGRATING FACTOR

$$\text{I.F.} = e^{\int \frac{4}{x} dx} = e^{4 \ln x} = e^{\ln x^4} = x^4$$

MULTIPLY THROUGH WE OBTAIN

$$x^4 \frac{dy}{dx} + x^4 \left( \frac{4y}{x} \right) = x^4 (6x - 5)$$

$$x^4 \frac{dy}{dx} + 4x^3 y = 6x^5 - 5x^4$$

$$\frac{d}{dx} (x^4 y) = 6x^5 - 5x^4$$

$$x^4 y = \int 6x^5 - 5x^4 dx$$

$$x^4 y = x^6 - x^5 + C$$

$$y = x^2 - x + \frac{C}{x^4}$$

APPLY CONDITION  $x=1, y=1$

$$1 = 1 - 1 + \frac{C}{1}$$

$$\therefore C = 1$$

THUS WE HAVE

$$y = x^2 - x + \frac{1}{x^4}$$

# IYGB - FP2 PAPER Q - QUESTION 2

PROCEED AS FOLLOWS

$$\int_0^e x^2 \ln x \, dx = \lim_{h \rightarrow 0^+} \left[ \int_h^e x^2 \ln x \, dx \right]$$

INTEGRATION BY PARTS

$$\dots = \lim_{h \rightarrow 0^+} \left[ \left[ \frac{1}{3} x^3 \ln |x| \right]_h^e - \int_h^e \frac{1}{3} x^2 \, dx \right]$$

$\ln x$	$\frac{1}{x}$
$\frac{1}{3} x^3$	$x^2$

$$= \lim_{h \rightarrow 0^+} \left[ \left[ \frac{1}{3} x^3 \ln |x| - \frac{1}{9} x^3 \right]_h^e \right]$$

$$= \lim_{h \rightarrow 0^+} \left[ \left( \frac{1}{3} e^3 \ln e - \frac{1}{9} e^3 \right) - \left( \frac{1}{3} h^3 \ln h - \frac{1}{9} h^3 \right) \right]$$

$$= \lim_{h \rightarrow 0^+} \left[ \frac{1}{3} e^3 - \frac{1}{9} e^3 - \frac{1}{3} h^3 \ln h + \frac{1}{9} h^3 \right]$$

NOW LOOKING AT THE TERM  $h^3 \ln h$ ,  $h^3$  TENDS TO ZERO "FASTER" THAN  $\ln h$  TENDS TO  $-\infty$ , I.E.  $\lim_{h \rightarrow 0^+} (h^3 \ln h) = 0$

$$\dots = \frac{1}{3} e^3 - \frac{1}{9} e^3 - 0 + 0$$

$$= \frac{2}{9} e^3$$

AS REQUIRED



## 1YGB - FP2 PAPER Q - QUESTION 3

START BY "SPLITTING THE FRACTION"

$$f(x) = \frac{e^x + 1}{2e^{\frac{1}{2}x}} = \frac{e^x}{2e^{\frac{1}{2}x}} + \frac{1}{2e^{\frac{1}{2}x}} = \frac{1}{2}e^{\frac{1}{2}x} + \frac{1}{2}e^{-\frac{1}{2}x}$$
$$= \cosh\left(\frac{1}{2}x\right)$$

Now  $\cosh u = 1 + \frac{u^2}{2!} + \frac{u^4}{4!} + \frac{u^6}{6!} + o(u^8)$

$$f(x) = 1 + \frac{\left(\frac{1}{2}x\right)^2}{2!} + \frac{\left(\frac{1}{2}x\right)^4}{4!} + \frac{\left(\frac{1}{2}x\right)^6}{6!} + o(x^8)$$

$$f(x) = 1 + \frac{1}{8}x^2 + \frac{1}{384}x^4 + \frac{1}{46080}x^6 + o(x^8)$$

### ALTERNATIVE USING EXPONENTIALS

- $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + o(x^4)$

- $e^{-x} = 1 - x + \frac{x^2}{2!} - \frac{x^3}{3!} + o(x^4)$

$$\therefore f(x) = \frac{1}{2}e^{\frac{1}{2}x} + \frac{1}{2}e^{-\frac{1}{2}x} = \frac{1}{2}\left(e^{\frac{1}{2}x} + e^{-\frac{1}{2}x}\right)$$

$$= \frac{1}{2} \left[ \begin{array}{l} 1 + \frac{1}{2}x + \frac{\frac{1}{4}x^2}{2!} + \frac{\frac{1}{8}x^3}{3!} + \frac{\frac{1}{16}x^4}{4!} + \frac{\frac{1}{32}x^5}{5!} + \frac{\frac{1}{64}x^6}{6!} + o(x^7) \\ 1 - \frac{1}{2}x + \frac{\frac{1}{4}x^2}{2!} - \frac{\frac{1}{8}x^3}{3!} + \frac{\frac{1}{16}x^4}{4!} - \frac{\frac{1}{32}x^5}{5!} + \frac{\frac{1}{64}x^6}{6!} + o(x^7) \end{array} \right]$$

$$= \frac{1}{2} \left[ 2 + \frac{1}{4}x^2 + \frac{1}{192}x^4 + \frac{1}{23040}x^6 + o(x^8) \right]$$

$$= 1 + \frac{1}{8}x^2 + \frac{1}{384}x^4 + \frac{1}{46080}x^6 + o(x^8)$$

AD 18506

# NYGB - FP2 PAPER Q - QUESTION 4

PROCEED BY IDENTITIES

$$\Rightarrow 2 \tanh^2 w = 1 + \operatorname{sech} w$$

$$\Rightarrow 2(1 - \operatorname{sech}^2 w) = 1 + \operatorname{sech} w$$

$$\Rightarrow 2 - 2 \operatorname{sech}^2 w = 1 + \operatorname{sech} w$$

$$\Rightarrow 0 = 2 \operatorname{sech}^2 w + \operatorname{sech} w - 1$$

$$\Rightarrow (2 \operatorname{sech} w - 1)(\operatorname{sech} w + 1)$$

$$\Rightarrow \operatorname{sech} w = \begin{cases} -1 \\ \frac{1}{2} \end{cases}$$


$$\Rightarrow \cosh w = \begin{cases} -2 \\ 2 \end{cases} \quad \leftarrow (\cosh w \geq 1)$$

$$\Rightarrow w = \pm \operatorname{arcosh} 2$$

$$\Rightarrow w = \pm \ln(2 + \sqrt{2^2 - 1})$$

$$\Rightarrow \underline{w = \pm \ln(2 + \sqrt{3})}$$

$$\begin{aligned} 1 + \tan^2 \theta &\equiv \sec^2 \theta \\ 1 - \tanh^2 \theta &\equiv \operatorname{sech}^2 \theta \\ 1 - \operatorname{sech}^2 \theta &\equiv \tanh^2 \theta \end{aligned}$$





- 1 -

## IYGB - FP2 PAPER Q - QUESTION 5

$$\frac{d^2y}{dx^2} - 3\frac{dy}{dx} + 2y = 10\sin x \quad y(0) = 6, \quad y'(0) = 5$$

AUXILIARY EQUATION

$$\lambda^2 - 3\lambda + 2 = 0$$

$$(\lambda - 2)(\lambda - 1) = 0$$

$$\lambda = \begin{matrix} 1 \\ 2 \end{matrix}$$

COMPLEMENTARY FUNCTION

$$y = Ae^x + Be^{2x}$$

PARTICULAR INTEGRAL BY INSPECTION

$$y = P\cos x + Q\sin x$$

$$y' = -P\sin x + Q\cos x$$

$$y'' = -P\cos x - Q\sin x$$

SUBSTITUTE INTO THE O.D.E

$$\Rightarrow (-P\cos x - Q\sin x) - 3(-P\sin x + Q\cos x) + 2(P\cos x + Q\sin x) \equiv 10\sin x$$

$$\Rightarrow \left. \begin{array}{l} -P\cos x - Q\sin x \\ -3Q\cos x + 3P\sin x \\ +2P\cos x + 2Q\sin x \end{array} \right\} \equiv 10\sin x$$

$$\Rightarrow (P - 3Q)\cos x + (3P + Q)\sin x \equiv 10\sin x$$

$$\begin{aligned} \bullet P - 3Q &= 0 \\ P &= 3Q \end{aligned}$$

$$\begin{aligned} \bullet 3P + Q &= 10 \\ 3(3Q) + Q &= 10 \\ 10Q &= 10 \\ \underline{Q} &= \underline{1} \end{aligned}$$

$$\& \quad \underline{P} = \underline{3}$$

# IYGB - FP2 PAPER Q - QUESTION 5

∴ PARTICULAR INTEGRAL

$$y = 3\cos x + \sin x$$

∴ GENERAL SOLUTION

$$y = Ae^{2x} + Be^{2x} + 3\cos x + \sin x$$

DIFFERENTIATE W.R.T  $x$  & APPLY CONDITIONS

$$\frac{dy}{dx} = Ae^{2x} + 2Be^{2x} - 3\sin x + \cos x$$

•  $x=0, y=6 \Rightarrow 6 = A + B + 3$

$\Rightarrow A + B = 3$

•  $x=0, \frac{dy}{dx}=5 \Rightarrow 5 = A + 2B + 1$

$\Rightarrow A + 2B = 4$

∴  $B=1 \quad A=2$

FINALLY WE OBTAIN

$y = 2e^{2x} + e^{2x} + 3\cos x + \sin x$



-|-

## (YGB - FP2 PAPER Q - QUESTION 6)

$$\text{USING } \frac{d}{dx} (\arctan f(x)) = \frac{1}{1+(f(x))^2} \times f'(x)$$

$$\frac{d}{dx} \left[ \arctan \left( \frac{x}{\sqrt{4-x^2}} \right) \right] = \frac{d}{dx} \left[ \arctan \left[ \frac{x}{(4-x^2)^{\frac{1}{2}}} \right] \right]$$

$$= \frac{1}{1 + \frac{x^2}{4-x^2}} \times \frac{(4-x^2)^{\frac{1}{2}} \times 1 - x \times \frac{1}{2}(4-x^2)^{-\frac{1}{2}}(-2x)}{(4-x^2)^1}$$

$$= \frac{1}{1 + \frac{x^2}{4-x^2}} \times \frac{(4-x^2)^{\frac{1}{2}} + x^2(4-x^2)^{-\frac{1}{2}}}{(4-x^2)}$$

$$= \frac{(4-x^2)^{\frac{1}{2}} + x^2(4-x^2)^{-\frac{1}{2}}}{(4-x^2) + x^2}$$

$$= \frac{(4-x^2)^{-\frac{1}{2}} \left[ (4-x^2)^1 + x^2 \right]}{(4-x^2) + x^2}$$

$$= \frac{1}{\sqrt{4-x^2}}$$

# YGB - FP2 PAPER Q - QUESTION 7

a) USING THE STANDARD FORMULA FOR THE AREA IN POLARS

$$\begin{aligned} \text{AREA} &= \int_{\theta_1}^{\theta_2} \frac{1}{2} r^2 d\theta = \int_0^{\frac{\pi}{2}} \frac{1}{2} (\sin 2\theta)^2 d\theta \\ &= \int_0^{\frac{\pi}{2}} \frac{1}{2} \sin^2 2\theta d\theta \end{aligned}$$

NOW USING THE TRIGONOMETRIC IDENTITY FOR COSINE DOUBLE ANGLE

$$\cos 2A \equiv 1 - 2\sin^2 A$$

$$\cos 4A \equiv \cos[2(2A)] \equiv 1 - 2\sin^2 2A$$

$$\sin^2 2A \equiv \frac{1}{2} - \frac{1}{2} \cos 4A$$

$$\begin{aligned} \text{AREA} &= \int_0^{\frac{\pi}{2}} \frac{1}{2} \left[ \frac{1}{2} - \frac{1}{2} \cos 4\theta \right] d\theta \\ &= \int_0^{\frac{\pi}{2}} \frac{1}{4} - \frac{1}{4} \cos 4\theta \\ &= \left[ \frac{1}{4} \theta - \frac{1}{16} \sin 4\theta \right]_0^{\frac{\pi}{2}} \\ &= \left( \frac{\pi}{2} \times \frac{1}{4} - 0 \right) - (0 - 0) \\ &= \frac{\pi}{8} \end{aligned}$$



YGB - FP2 PAPER Q - QUESTION 7

b) For "HORIZONTAL TANGENT"  $\frac{dy}{dx} = 0$

$$\frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = 0$$

$$\frac{dy}{d\theta} = \frac{d}{d\theta}(r \sin \theta) = \frac{d}{d\theta}(\sin 2\theta \sin \theta) = 0$$

DIFFERENTIATE & SOLVE THE EQUATION

$$\Rightarrow 2\cos 2\theta \sin \theta + \sin 2\theta \cos \theta = 0$$


$$\Rightarrow 2\sin \theta (2\cos^2 \theta - 1) + 2\sin \theta \cos^2 \theta = 0$$

$$\Rightarrow 2\sin \theta [3\cos^2 \theta - 1] = 0$$


$$\therefore \cancel{\sin \theta = 0} \quad \cos \theta = \frac{1}{\sqrt{3}} \quad \cos \theta = -\frac{1}{\sqrt{3}}$$

$$\therefore \theta = \arccos\left(\frac{1}{\sqrt{3}}\right)$$

$$\begin{aligned} \therefore r &= \sin 2\theta = 2\sin \theta \cos \theta \\ &= 2 \times \frac{\sqrt{2}}{\sqrt{3}} \times \frac{1}{\sqrt{3}} = \frac{2}{3}\sqrt{2} \end{aligned}$$



Polar coordinates of P  $\left(\frac{2}{3}\sqrt{2}, \arccos\frac{1}{\sqrt{3}}\right)$



$$x = r \cos \theta = \frac{2}{3}\sqrt{2} \times \frac{1}{\sqrt{3}} = \frac{2\sqrt{2}}{3\sqrt{3}} = \frac{2}{9}\sqrt{6}$$

$$y = r \sin \theta = \frac{2}{3}\sqrt{2} \left(\frac{\sqrt{2}}{\sqrt{3}}\right) = \frac{4}{3\sqrt{3}} = \frac{4}{9}\sqrt{3}$$

CARTESIAN COORDINATES  $\left(\frac{2}{9}\sqrt{6}, \frac{4}{9}\sqrt{3}\right)$



# 1YGB - FP2 PAPER Q - QUESTION 8

USING THE BINOMIAL

$$(a-1)^3 = a^3 - 3a^2 \cdot 1 + 3a \cdot 1^2 - 1^3$$

$$(a-1)^3 = a^3 - 3a^2 + 3a - 1$$

COMPARING WE OBTAIN

$$\Rightarrow z^3 - 3z^2 + 3z - 65 = 0$$

$$\Rightarrow \underbrace{z^3 - 3z^2 + 3z - 1}_{(z-1)^3} - 64 = 0$$

$$(z-1)^3 = 64$$

USING EXPONENTIALS

$$\Rightarrow (z-1)^3 = 64 e^{0i + 2k\pi i} \quad k = 0, 1, 2$$

$$\Rightarrow (z-1)^3 = 64 e^{2k\pi i}$$

$$\Rightarrow z-1 = 64^{\frac{1}{3}} e^{\frac{2k\pi i}{3}}$$

$$\Rightarrow z = 1 + 4e^{\frac{2k\pi i}{3}}$$

•  $z_0 = 1 + 4e^0 = 1 + 4 = 5$

•  $z_1 = 1 + 4e^{\frac{2\pi i}{3}} = 1 + 4\left(\cos\frac{2\pi}{3} + i\sin\frac{2\pi}{3}\right) = 1 + 4\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)$   
 $= -1 + 2\sqrt{3}i$

•  $z_2 = 1 + 4e^{\frac{4\pi i}{3}} = 1 + 4\left(\cos\frac{4\pi}{3} + i\sin\frac{4\pi}{3}\right) = 1 + 4\left(-\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)$   
 $= -1 - 2\sqrt{3}i$

$\therefore z = \begin{cases} 5 \\ -1 + 2\sqrt{3}i \\ -1 - 2\sqrt{3}i \end{cases}$



1XGB - FP2 PART Q - QUESTION 9

a)  $f(r) = \frac{1}{(r+1)(r-1)} = \frac{\frac{1}{2}}{r-1} - \frac{\frac{1}{2}}{r+1} = \frac{1}{2(r-1)} - \frac{1}{2(r+1)}$   
 "BY COVER UP" (INSPECTION)

b) USING PART (a) "DOUBLED" FOR SIMPLICITY

$$\frac{2}{r^2-1} \equiv \frac{2}{(r+1)(r-1)} \equiv \frac{1}{r-1} - \frac{1}{r+1}$$

• $r=2$	$\frac{2}{2^2-1} = \frac{1}{1} - \frac{1}{3}$
• $r=3$	$\frac{2}{3^2-1} = \frac{1}{2} - \frac{1}{4}$
• $r=4$	$\frac{2}{4^2-1} = \frac{1}{3} - \frac{1}{5}$
• $r=5$	$\frac{2}{5^2-1} = \frac{1}{4} - \frac{1}{6}$
• $r=6$	$\frac{2}{6^2-1} = \frac{1}{5} - \frac{1}{7}$
• $\vdots$	$\vdots$
• $r=n-1$	$\frac{2}{(n-1)^2-1} = \frac{1}{n-2} - \frac{1}{n}$
• $r=n$	$\frac{2}{n^2-1} = \frac{1}{n-1} - \frac{1}{n+1}$

$$\sum_{r=2}^n \frac{2}{r^2-1} = 1 + \frac{1}{2} - \frac{1}{n} - \frac{1}{n+1}$$

$$2 \sum_{r=2}^n \frac{1}{r^2-1} = \frac{3}{2} - \frac{1}{n} - \frac{1}{n+1}$$

$$\sum_{r=2}^n \frac{1}{r^2-1} = \frac{3}{4} - \frac{1}{2n} - \frac{1}{2(n+1)}$$

c) AS  $n \rightarrow \infty$  THE SUM TEND TO  $\frac{3}{4}$